



CHCP-I-T Cryopreserved Plateable Cynomolgus Hepatocytes for Induction and Transporter assays		
Cell Specification		
Lot CH140120		Batch Release: June 9, 2017
Species: <i>Macaca fascicularis</i> Gender: female Age: 3 years 10 months		Serology: negative for Filovirus/Ebola-like, SRV, SIV, STLV-1
Cryopreservation: Date: January 20, 2014 Amount per vial: 10.2 x 10 ⁶ cells		Thawing: Post-thaw viability: 89 % Post-thaw yield per vial: 5.1 x 10 ⁶ cells Recovery: 50 %
Phase contrast on day 1 after thawing 		Phase contrast on day 3 after thawing 
Recommended seeding density on collagen-coated plates: 24well plate – 300,000 cells/well 96well plate – 70,000 cells/well Culture in Human Hepatocyte Maintenance Medium (HHMM).		
CYP P450 activity in culture after thawing:	pmol/(mg × min)	X-fold induction
Ethoxyresorufin-O-deethylation:	24well: 65.6 ± 0.3	19.9
Induction with 25 µM beta-naphthoflavone	96well: 133.0 ± 18.1	19.7

Uptake Transporters: uptake of 10 μ M Estrone 3-sulfate (E_3S) with or without competitive inhibitor Bromosulfophthalein (BSP, 100 μ M) in cryopreserved hepatocytes after 2 min incubation.		
Activity of uptake transporters in culture after thawing	Intracellular E_3S (pmol/mg \times min)	Inhibition (%)
Without BSP	371 \pm 102	
With BSP	260 \pm 72	29.9
Efflux Transporter: after 5 min preincubation with 10 μ M Talinolol (Tal), P-glycoprotein (P-gp) mediated Efflux of Tal with or without the specific P-gp inhibitor PSC833 (10 μ M) in cryopreserved hepatocytes after 60 min incubation was analyzed.		
Activity of P-gp in culture after thawing	Intracellular Tal (pmol/mg)	
Without PSC833	177 \pm 30	
With PSC833	234 \pm 58	
<p>Note: Yield, viability, recovery and activity assays were performed at PRIMACYT using PRIMACYT's manual for thawing, plating and culture of primary cryopreserved hepatocytes.</p> <p>Issued by: A. Ullrich Checked by: C. Garve</p>		