

Thawing, Plating and Culture of Primary Cryopreserved Hepatocytes

Updated 21.11.2018

Version 06

Required and recommended media and consumables

- Thawing and Plating Kit consists of
 - HTM: Hepatocyte Thawing Medium
 - HWM: Hepatocyte Washing Medium
 - HPM-cryo: Hepatocyte Plating Medium for cryopreserved hepatocytes
- For the use of cryopreserved hepatocytes in suspension only HTM and HWM are required. Both components are not sold separately, only in combination with the kit.
- HHMM (Human Hepatocyte Maintenance Medium): for serum-free culture of human, monkey, and dog hepatocytes (not included in this kit)
- Collagen coated cell culture plates (not included in this kit)

1. Arrival of the cryopreserved cells in your laboratory

- Place the cryogenic vial with frozen hepatocytes immediately into the gas phase of liquid nitrogen tank or store at/below -150 °C

2. Thawing and Plating of primary hepatocytes

- Warm water bath, HTM, and HWM to 37 °C
- Set HPM-cryo to room temperature
- Remove the vial with hepatocytes from liquid nitrogen/-150 °C and place it immediately into the 37 °C warm water bath until the cell suspension is thawed (approx. 1-2 min)
- Spray 70 % ethanol on the cryogenic vial for disinfection
- Transfer the cell suspension into the tube with HTM
- Wash the cryogenic vial with 0.5-1 ml HWM to remove the cells completely and combine it with the cells in the tube
- Add HWM to a final volume of 50 ml
- Rotate the tube slowly two or three times
- Pellet the hepatocytes by centrifugation at 100 x g and 20 °C for 10 min
- Remove the supernatant, gently loosen the cells without any additional medium by gently agitating the bottom of the tube. Do not vortex or shake the cells.
- Wash the loosen cells with 20 ml HWM followed by centrifugation at 50 x g and 20 °C for 5 min
- Remove the supernatant, gently loosen the cells without any additional medium by gently agitating the bottom of the tube. Do not vortex or shake the cells.
- Re-suspend the pellet in an appropriate volume of HPM-cryo; 2-4 ml of HPM-cryo is recommended when counting one cryogenic vial with hepatocytes, 10 mio./vial.

- Determine cell viability and live cell number with the trypan blue exclusion test in a counting chamber (do not use automated cell counter)
- Seed the cells at a density recommended in the corresponding data sheet of each individual lot.
- Adjust cell suspension to the desired density for plating with HPM-cryo

Note: the actual seeding density may vary from lot to lot. The recommended seeding density for each lot is stated on the accompanying data sheet.

- Recommended volume of plating medium per well:
 - 6well: 1-2 ml
 - 12well: 1 ml
 - 24well: 0.5 ml
 - 96well: 100 µl
- Let the cells attach for at least 6-7 h at 37 °C and 5 % CO₂, do not let the cells attach overnight

3. Culture of primary hepatocytes

- After attachment of cells: change medium
- Wash the cells with warm PBS (1-2 times)
 - 6well: 2 ml/well
 - 12well: 1 ml/well
 - 24well: 0.5 ml/well
 - 96well: 50 µl/well
- Heat the culture medium to 37 °C (no longer than 15 min)
- Replace the Hepatocyte Plating Medium with Human Hepatocyte Maintenance Medium (HHMM)
 - 6well: 1 ml/well
 - 12well: 0.5 ml/well
 - 24well: 0.3 ml/well
 - 96well: 50 µl/well
- Change the medium daily (especially when plated at high cell density)
- Change the medium quickly, do not let the cells dry

FOR IN VITRO RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC PROCEDURES.